

Radiative corrections and parity nonconservation in heavy atoms

A. I. Milstein*

Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia

O. P. Sushkov†

School of Physics, University of New South Wales, Sydney 2052, Australia

I. S. Terekhov‡

Novosibirsk University, 630090 Novosibirsk, Russia

(Dated: February 1, 2008)

The self-energy and the vertex radiative corrections to the effect of parity nonconservation in heavy atoms are calculated analytically in orders $Z\alpha^2$ and $Z^2\alpha^3\ln(\lambda_C/r_0)$, where λ_C and r_0 being the Compton wavelength and the nuclear radius, respectively. The value of the radiative correction is -0.85% for Cs and -1.41% for Tl. Using these results we have performed analysis of the experimental data on atomic parity nonconservation. The obtained values of the nuclear weak charge, $Q_W = -72.90(28)_{exp}(35)_{theor}$ for Cs, and $Q_W = -116.7(1.2)_{exp}(3.4)_{theor}$ for Tl, agree with predictions of the standard model. As an application of our approach we have also calculated analytically dependence of the Lamb shift on the finite nuclear size.

PACS numbers: 11.30.Er, 31.30.Jv, 32.80.Ys

Atomic parity nonconservation (PNC) has now been measured in bismuth [1], lead [2], thallium [3], and cesium [4]. Analysis of the data provides an important test of the standard electroweak model and imposes constraints on new physics beyond the model, see Ref. [5]. The analysis is based on the atomic many-body calculations for Tl, Pb, and Bi [6] and for Cs [7, 8] (see also more recent Refs. [9, 10]). Both the experimental and the theoretical accuracy is best for Cs. Therefore, this atom provides the most important information on the standard model in the low-energy sector. The analysis performed in Ref. [4] has indicated a deviation of the measured weak charge value from that predicted by the standard model by 2.5 standard deviations σ .

In the many-body calculations [6, 7, 8] the Coulomb interaction between electrons was taken into account, while the magnetic interaction was neglected. The contribution of the magnetic (Breit) electron-electron interaction was calculated in the papers [11, 12]. It proved to be much larger than a naive estimate, and it shifted the theoretical prediction for PNC in Cs.

Radiative correction to the nuclear weak charge due to renormalization from the scale of the W-boson mass down to zero momentum has been calculated long time ago, see Refs. [13, 14]. This correction is always accounted in the analysis of data. However, another important class of radiative corrections was lost in the analysis of atomic PNC. This fact has been pointed out in the recent work [15] that demonstrated that there are corrections $\sim Z\alpha^2$ caused by the strong electric field of the nucleus. Here Z is the nuclear charge and α is the fine structure constant. The simplest correction of this type is due to the Uehling potential. It has been calculated numerically in Ref. [16] and analytically in our paper

[17]. In the paper [17] we have also analyzed the general structure of the strong field radiative corrections. It has been shown that except of the usual perturbative parameter $Z\alpha$ there is an additional parameter $\ln(\lambda_C/r_0)$ where λ_C is the electron Compton wavelength and r_0 is the nuclear radius.

In this letter we present the results of calculations of the strong field radiative corrections to the atomic PNC effect in orders $Z\alpha^2$ and $Z^2\alpha^3\ln(\lambda_C/r_0)$. Details of calculations will be presented elsewhere [18]. Using our results we reanalyze the experimental data. Agreement with the standard model is excellent. As an application of our approach we have also calculated the dependence of the Lamb shift on the finite nuclear size. Agreement of our analytical formula with results of previous computations [19, 20] is perfect.

The strong relativistic enhancement makes PNC radiative corrections different from previously considered radiative corrections to the hyperfine structure. The relativistic enhancement factor is proportional to $R \sim (\lambda_C/(Z\alpha r_0))^{2(1-\gamma)}$, where $\gamma = \sqrt{1 - (Z\alpha)^2}$. The factor is $R \approx 3$ for Cs and $R \approx 9$ for Tl, Pb, and Bi [21]. The logarithmic enhancement of radiative corrections mentioned above is closely related to the existence of the factor R . The Feynman diagram for the leading contribution to the PNC matrix element between $p_{1/2}$ and $s_{1/2}$ states as well as diagrams for radiative corrections are shown in Fig.1(a) and Fig.1(b-f), respectively. The diagram Fig.1(b) corresponds to a modification of the electron wave function because of the vacuum polarization. This correction calculated analytically in Ref. [17] reads

$$\delta_b = \alpha \left(\frac{1}{4} Z\alpha + \frac{2(Z\alpha)^2}{3\pi\gamma} [\ln^2(b\lambda_C/r_0) + f] \right), \quad (1)$$




FIG. 1: (a) Leading contribution to the PNC matrix element. (b-f) Radiative corrections. The double line is the exact electron Green's function in the Coulomb field of the nucleus, the cross denotes the nucleus, the zigzag and the dashed lines denote Z-boson and photon, respectively.

where $b = \exp(1/(2\gamma) - C - 5/6)$, $C \approx 0.577$ is the Euler constant, and $f \sim 1$ is some smooth function of $Z\alpha$ independent of r_0 . Hereafter we denote by δ the relative value of the correction. So, Eq. (1) represents the ratio of diagrams Fig.1(b) and Fig.1(a).

The renormalization of the nuclear weak charge Q_W from the scale of the W-boson mass down to $q = 0$ was performed in Refs. [13, 14]. However, as it has been pointed out in Ref. [17], atomic experiments correspond to $q \sim 1/r_0 \sim 30\text{MeV}$. The correction due to renormalization from $q = 0$ to $q = 1/r_0$ is described by diagrams Fig.1(c) and Fig.1(d). It has the form [17]

$$\delta_{cd} = \frac{4\alpha Z}{3\pi Q_W} (1 - 4\sin^2\theta_W) \ln(\lambda_C/r_0) \sim -0.1\%, \quad (2)$$

where θ_W is the Weinberg angle, $\sin^2\theta_W \approx 0.2230$, see Ref. [5].

Diagrams Fig.1(e) and Fig.1(f) correspond to the contributions of the electron self-energy operator and the vertex operator, respectively. Each of these diagrams is not invariant with respect to the gauge transformation of the electromagnetic field while their sum is gauge invariant. It has been demonstrated in Ref. [17] that the correction $\delta_e + \delta_f$ is of the form

$$\delta_{ef} = \delta_e + \delta_f = A \ln(b\lambda_C/r_0) + B, \quad (3)$$

where A and B are some functions of $Z\alpha$, and the constant b is defined after Eq.(1) In the present paper we derive the functions A and B analytically in the leading approximation in the parameter $Z\alpha$, when $A \propto \alpha(Z\alpha)^2$ and $B \propto \alpha(Z\alpha)$.

The simplest part of the work is calculation of the $\alpha(Z\alpha)$ contribution to B . It is convenient to use the Fried-Yennie gauge [22] together with the effective operators approach [23] where the corrections under discussion coincide with that for the forward scattering amplitude, see Fig.2. The result of our calculation for the non-logarithmic term in Eq.(3) reads

$$B = -\alpha(Z\alpha) \left(\frac{7}{12} + 2 \ln 2 \right). \quad (4)$$




FIG. 2: $\alpha(Z\alpha)$ self-energy and vertex radiative corrections. The zigzag line denotes Z-boson, and the dashed line denotes photon.

For calculation of the function A (the logarithmic part in Eq. (3)) we have used the Feynman gauge. There are two contributions to A , the self-energy contribution, A_{SE} , given by the diagram Fig.1(e), and the vertex contribution A_V given by Fig.1(f). It is convenient to represent the self-energy operator as a series in powers of the Coulomb field of the nucleus, $\Sigma = \Sigma_0 + \Sigma_1 + \Sigma_2 + \dots$, see Fig.3, and perform calculations in the momentum space. Distances $r_0 \ll r \ll \lambda_C$ where the logarithmic




FIG. 3: The electron self energy expanded in powers of the Coulomb field. The solid line is the free electron Green's function, the cross denotes the nucleus, and the dashed line denotes the photon.

terms come from correspond to momenta $m \ll p \ll 1/r_0$, where m is the electron mass. Therefore, the mass and the electron binding energy can be neglected in electron propagators. It is necessary to make the ultraviolet regularization of the operators Σ_0 and Σ_1 . Each of the operators depends on the parameter of the regularization. However, the contribution of $\Sigma_{01} = \Sigma_0 + \Sigma_1$ to A_{SE} is independent of the regularization parameter because of the Ward identity. The operator Σ_2 does not require any regularization. Our results for contributions of Σ_{01} and Σ_2 to A_{SE} read

$$\begin{aligned} A_{01} &= -\frac{\alpha(Z\alpha)}{\pi}, \\ A_2 &= -\frac{\alpha(Z\alpha)}{6\pi}(\pi^2 - 9), \\ A_{SE} &= A_{01} + A_2 = -\frac{\alpha(Z\alpha)^2}{6\pi}(\pi^2 - 3). \end{aligned} \quad (5)$$

Next is the logarithmic contribution of the vertex operator described by Fig.1(f). The coordinate representation is the most convenient for this part of the problem, and we use this representation. We have also used an integral representation of the Green function derived in Ref.[24] (see also Ref. [25]). Keeping in mind the contact nature of the PNC interaction and the logarithmic accuracy of the calculation, one can demonstrate that only the angular momentum $j = 1/2$ is significant in the partial expansion of the electron Green's function in the external Coulomb field. A part of the vertex independent of

$Z\alpha$ requires an ultraviolet regularization and hence is dependent of the regularization parameter. However, due to the Ward identity, the contribution corresponding to this part cancels out exactly the renormalization of the wave function, see e.g. Ref. [26]. Another part is dependent on $Z\alpha$ and gives the following contribution to the logarithmic term in Eq. (3)

$$A_V = -\frac{\alpha(Z\alpha)^2}{\pi} \left(\frac{17}{4} - \frac{\pi^2}{3} \right) . \quad (6)$$

Together with Eq. (5) this gives the final result for A

$$A = -\frac{\alpha(Z\alpha)^2}{\pi} \left(\frac{15}{4} - \frac{\pi^2}{6} \right) . \quad (7)$$

Thus, according to Eqs. (3), (4), and (7) the total relative correction to the PNC matrix element due to the self-energy and the vertex operators reads

$$\begin{aligned} \delta_{ef} = & -\alpha \left[(Z\alpha) \left(\frac{7}{12} + 2 \ln 2 \right) \right. \\ & \left. + \frac{(Z\alpha)^2}{\pi} \left(\frac{15}{4} - \frac{\pi^2}{6} \right) \ln(b\lambda_C/r_0) \right] . \quad (8) \end{aligned}$$

This correction is plotted in Fig. 4 versus the nuclear charge Z (the long-dashed line). The solid line in the same figure shows the total radiative correction to the PNC effect that includes both δ_{ef} (Eq. (8) and δ_b (Eq. (1)).

The radiative shift of the atomic energy levels (Lamb shift) depends on the finite nuclear size. This correction has a very similar structure to that of the PNC radiative correction since the effective sizes of the perturbation sources in both cases are much smaller than λ_C . The self-energy and the vertex corrections to the finite-nuclear-size effect (SEVFNS) for $s_{1/2}$ -state have been calculated earlier analytically in order $\alpha(Z\alpha)$ in Refs.[27, 28]. The corrections for $1s_{1/2}$ -, $2s_{1/2}$ -, and $2p_{1/2}$ -states have been calculated numerically exactly in $Z\alpha$ in Refs.[19, 30, 31]. The structure of higher in $Z\alpha$ corrections and their logarithmic dependence on the nuclear size has not been understood. We have applied our approach to the SEVFNS problem and found the following expression for the $s_{1/2}$ -state relative correction

$$\begin{aligned} \Delta_s = & -\alpha \left[(Z\alpha) \left(\frac{23}{4} - 4 \ln 2 \right) \right. \\ & \left. + \frac{(Z\alpha)^2}{\pi} \left(\frac{15}{4} - \frac{\pi^2}{6} \right) \ln(b\lambda_C/r_0) \right] . \quad (9) \end{aligned}$$

Linear in $Z\alpha$ term agrees with results of Refs. [27, 28]. The logarithmic term coincides with that in Eq. (8) for the PNC correction. Moreover, the logarithmic term in the SEVFNS correction Δ_p for $p_{1/2}$ state is also equal to that in Eqs.(8) and (9). The reason for this equality is very simple. The logarithmic terms come from small

distances ($r \ll \lambda_C$) where the electron mass can be neglected. When the mass is neglected the relative matrix elements for the PNC radiative correction and for SEVFNS are equal. The correction Δ_s given by Eq. (9) is shown in Fig. 4 by the dashed line. Results of the computations [19, 20] for $1s$ and $2s$ states are shown by circles and diamonds, respectively. The agreement is perfect. It




FIG. 4: Relative radiative corrections (%) for the PNC and for the finite-nuclear-size effects versus the nuclear charge Z . The long-dashed line shows the correction δ_{ef} given by Eq.(8) (Fig.1(e,f)). The solid line shows the total radiative correction to the PNC effect that includes both δ_{ef} and δ_b (Eq.(1), Fig.1(b)). The dashed line shows the correction Δ_s given by Eq.(9). Results of computations of Δ_s for $1s$ and $2s$ states [19, 20] are shown by circles and diamonds, respectively.

is interesting to find also the $\alpha(Z\alpha)$ -term in Δ_p . Using our result for the logarithmic term in Δ_p and results of numerical calculations of this correction for $2p_{1/2}$ -state, Ref. [19, 20], we find that the coefficient in $\alpha(Z\alpha)$ -term is -1.9 .

It has been recently suggested in Ref. [29] that the following "precise relation" $\delta_{ef} = (\Delta_s + \Delta_p)/2$ is valid. Values of the corrections obtained in our work do not agree with this relation. The "derivation" in Ref. [29] is based on the obviously wrong assumption that there is a gauge in which the vertex contributions to δ_{ef} , Δ_s , and Δ_p vanish simultaneously. Although it is possible to vanish the vertex correction to each of these quantities by choosing an appropriate gauge, these are three different gauges. This is why the above relation is wrong.

Now we can perform a consistent analysis of the experimental data on atomic parity violation since all the contributions are known. In our analysis for Cs we have included the theoretical value of the PNC amplitude from Refs. [7, 8, 9, 10] as well as -0.61% correction due to the Breit interaction [12], -0.42% radiative correction calculated in the present work, -0.2% neutron skin correction [32], -0.08% correction due to the renormalization of Q_W from the atomic momentum transfer $q \sim 30\text{MeV}$ down to $q = 0$ [17], and $+0.04\%$ contribution from the electron-

electron weak interaction [17]. Using these theoretical results we obtain from the data [4] the following value of the nuclear weak charge Q_W at zero momentum transfer

$$Cs: \quad Q_W = -72.90 \pm (0.28)_{ex} \pm (0.35)_{theor}. \quad (10)$$

This value agrees with prediction of the standard model, $Q_W = -73.09 \pm 0.03$, see Ref. [5]. We have used the neutron skin correction in our analysis. However, in our opinion, status of the correction is not quite clear because data on the neutron distribution used in Ref. [32] are not quite consistent with the data on neutron distributions obtained from proton scattering, see e. g. Ref. [33].

In the analysis for Tl we have included the theoretical value of the PNC amplitude from Refs. [6], as well as -0.88% correction due to the Breit interaction [34], -0.51% radiative correction calculated in the present work, -0.2% neutron skin correction, -0.08% correction due to the renormalization of Q_W from the atomic momentum transfer $q \sim 30\text{MeV}$ down to $q = 0$ [17], and $+0.01\%$ contribution from the electron-electron weak interaction [17]. Using these theoretical results we obtain from the data [3] the following value of the nuclear weak charge Q_W at zero momentum transfer

$$Tl: \quad Q_W = -116.7 \pm (1.2)_{ex} \pm (3.4)_{theor}. \quad (11)$$

This agrees with prediction of the standard model, $Q_W = -116.7 \pm 0.1$, see Ref. [5].

Concluding, we have calculated analytically for the first time all the strong electric field radiative corrections to the effect of atomic parity violation. This calculation has allowed us to perform a consistent analysis of the experimental data. Agreement with the standard model is within 0.5σ .

A.I.M gratefully acknowledge School of Physics at the University of New South Wales for warm hospitality and financial support during a visit.

* Email: A.I.Milstein@inp.nsk.su

† Email: sushkov@phys.unsw.edu.au

‡ Email: I.S.Terekhov@inp.nsk.su

- [1] L. M. Barkov and M. S. Zolotarev, JETP Lett., **27**, 357 (1978); M. J. D. MacPherson *et al.*, Phys. Rev. Lett., **67**, 2784 (1991).
- [2] D. M. Meekhof *et al.*, Phys. Rev. Lett., **71**, 3442 (1993).
- [3] N. H. Edwards *et al.*, Phys. Rev. Lett., **74**, 2654 (1995); P. A. Vetter *et al.*, Phys. Rev. Lett., **74**, 2658 (1995).

- [4] C. S. Wood *et al.*, Science **275**, 1759 (1997); S. C. Bennett and C. E. Wieman, Phys. Rev. Lett. **82**, 2484 (1999).
- [5] D. E. Groom *et al* Euro. Phys. J. C **15**, 1 (2000).
- [6] V. A. Dzuba, V. V. Flambaum, P. G. Silvestrov, and O. P. Sushkov, J. Phys. B **20**, 3297 (1987); Europhys. Lett. **7**, 413 (1988).
- [7] V. A. Dzuba, V. V. Flambaum, and O. P. Sushkov, Phys. Lett. **141A**, 147 (1989).
- [8] S. A. Blundell, J. Sapirstein, and W. R. Johnson, Phys. Rev. Lett. **65**, 1411 (1990); Phys. Rev. **D45**, 1602 (1992).
- [9] M. G. Kozlov, S. G. Porsev, and I. I. Tupitsyn, Phys. Rev. Lett. **86**, 3260 (2001).
- [10] V. A. Dzuba, V. V. Flambaum and J. S. M. Gingers, hep-ph/0204131.
- [11] A. Derevianko, Phys. Rev. Lett. **85**, 1618 (2000).
- [12] V. A. Dzuba, C. Harabati, W. R. Johnson, and M. S. Safronova, Phys. Rev. A **63**, 044103 (2001).
- [13] W. J. Marciano and A. Sirlin, Phys. Rev. D **27**, 552 (1983).
- [14] W. J. Marciano and J. L. Rosner, Phys. Rev. Lett. **65**, 2963 (1990).
- [15] O. P. Sushkov, Phys. Rev. A **63**, 042504 (2001);
- [16] W. R. Johnson, I. Bednyakov and G. Soff, Phys. Rev. Lett. **87**, 233001 (2001).
- [17] A. I. Milstein and O. P. Sushkov, Phys. Rev. A **66**, 022108 (2002).
- [18] A. I. Milstein, O. P. Sushkov and I. S. Terekhov, to be published.
- [19] K. T. Cheng, W. R. Johnson, and J. Sapirstein, Phys. Rev. A **47**, 1817 (1993).
- [20] W. R. Johnson and G. Soff, At. Data Nuc. Data Tables **33**, 405 (1985).
- [21] *Parity nonconservation in atomic phenomena*, I. B. Khriplovich, Gordon and Breach, 1991.
- [22] H. M. Fried and D. R. Yennie, Phys. Rev. **112**, 1391 (1958).
- [23] M. I. Eides, H. Grotch, and V. A. Shelyuto, Phys. Rep. **342**, 63 (2001).
- [24] A.I. Milstein, V.M. Strakhovenko, Phys. Lett. A **90**, 447 (1982).
- [25] R. N. Lee and A. I. Milstein, Phys. Lett. A **189**, 72 (1994).
- [26] K. Pachucki, Phys. Rev. A **54**, 1994 (1996).
- [27] K. Pachucki, Phys. Rev. A **48**, 120 (1993).
- [28] M. I. Eides, H. Grotch, Phys. Rev. A **56**, R2507 (1997).
- [29] M. Yu. Kuchiev and V. V. Flambaum, hep-ph/0206124.
- [30] S. A. Blundell, Phys. Rev. A **46**, 3762 (1992).
- [31] I. Lindgren, H. Persson, S. Salomonson, A. Ynnerman, Phys. Rev. A **47**, 4555 (1993).
- [32] A. Derevianko, Phys. Rev. A **65**, 012106 (2002);
- [33] G. D. Alkhazov *et al.*, Sov. J. Nucl. Phys. **26**, 357 (1977).
- [34] We found this value by rescaling the Breit correction for Cs [12], see discussion in Refs. [15, 17].